

# A STAND-REPLACEMENT PRESCRIBED BURN IN SAND PINE SCRUB

Kenneth W. Outcalt

U.S. Department of Agriculture, Forest Service, Southern Research Station, Athens, GA 30602

Cathryn H. Greenberg

U.S. Department of Agriculture, Forest Service, Bent Creek Experimental Forest, Asheville, NC 28806

## ABSTRACT

This paper describes fire characteristics and the immediate effects of a prescribed, high-intensity burn on a 12.2 hectare portion of a stand of Ocala sand pine scrub. The fire team on the Seminole District, Ocala National Forest used the BEHAVE fire model to predict the conditions needed to accomplish a stand-replacement burn. Suitable conditions arose on May 11, 1993 with temperature 26° Celsius, relative humidity 50%, wind 3 kilometers per hour, and fuel moisture (1-hour fuel) 7%. The area was burned by establishing a 40 meter blackline with backfires and then setting a headfire. The prolific smoke produced was not a problem due to selection of proper atmospheric conditions. Fire intensity was variable and affected by position and fuel loading, but on average was twice as high in the interior compared to the edge. The crowns of nearly all sand pine were severely scorched, and all trees subsequently died. Twenty-seven percent of the preburn snags were felled by the fire. Fire eliminated the shrub layer and reduced the litter layer thickness by 50%. The amount of bare ground was 0.1% in control plots and 17% in the burned area. Following the burn, light increased from 6 to 17% at ground level but only from 16 to 22% at breast height, because most needles remained on the overstory sand pine.

*Citation:* Outcalt, Kenneth W., and Cathryn H. Greenberg. 1998. A stand-replacement prescribed burn in sand pine scrub. Pages 141–145 in Teresa L. Pruden and Leonard A. Brennan (eds.). *Fire in ecosystem management: shifting the paradigm from suppression to prescription*. Tall Timbers Fire Ecology Conference Proceedings, No. 20. Tall Timbers Research Station, Tallahassee, FL.

## INTRODUCTION

The sand pine scrub community is a unique habitat with a suite of species that occur in no other community (Christman and Judd 1990). Mature scrub has an overstory of even-aged sand pine (*Pinus clausa* [Chapm. ex Engelm.] Vassey ex Sarg.) and a variably thick understory of sclerophyllous evergreen shrubs (Myers 1990). Typical understory species include myrtle oak (*Quercus myrtifolia*), sand live oak (*Q. geminata*), Chapman's oak (*Q. chapmanii*), turkey oak (*Q. laevis*), rusty lyonia (*Lyonia ferruginea*), rosemary (*Ceratiola ericoides*), scrub palmetto (*Sabal etonia*), and saw palmetto (*Serenoa repens*). Species richness and diversity of herbaceous plants are significantly less in mature stands compared to recently disturbed areas (Greenberg et al. 1995b). The sparse cover of herbs and grasses that do occur in mature scrub habitats typically include beak rush (*Rhynchospora megalocarpa*), milk peas (*Galactia* spp.), and broomsedge (*Andropogon* spp.). Lichens (*Cladonia* spp.) form extensive patches on the forest floor (Greenberg et al. 1995b).

Scrub dominated by the Ocala variety of sand pine (*Pinus clausa* var. *clausa* D.B. Ward) is native to the central ridge of Florida (Figure 1). It is also found on a strip of old dunes stretching from St. John's County south to the northern portion of Dade County on the east coast and from near Cedar Key south to Naples on the west coast (Small 1921, Harper 1927, Myers 1990). The largest concentration is the interior scrub, which occupies about 101,215 hectares on the Ocala National Forest (Brendemuehl 1990).

Sand pine scrub grows on deep droughty infertile sands of marine and aeolian origin. Water and wind formed these features as sea levels fluctuated during past glacial and interglacial periods (Kurz 1942, Laessle 1958, Brooks 1972). Because of washing and sorting during transport and deposition, soil parent material was nearly pure quartz sand (Laessle 1958). These processes produced soils that are almost exclusively Entisols and mostly Quartzipsamments (Myers 1990) typified by the Astatula, Lakeland, Paola, and St. Lucie soil series.

Because ground cover is sparse, Ocala sand pine scrub will not burn most of the time. Periodically, every 10 to 100 years, usually during a spring drought, high wind and temperature results in a catastrophic fire. Such fires kill the sand pine overstory and burn off the understory (Myers 1990). Heat from intense fire also opens many serotinous cones in crowns of sand pine, which releases seed for establishment of the next stand. Herbaceous species, including many endemics, also respond rapidly to the disturbance by invading, resprouting, or germinating from the soil seed bank (Johnson 1982, Hartnett and Richardson 1989).

Without high-intensity disturbance, the sand pine scrub would succeed to xeric oak hammock (Laessle 1958, Veno 1976) and lose many plant and animal species requiring young, open scrub sites (Greenberg et al. 1994, 1995a, 1995b). Most of the sand pine scrub on the Ocala National Forest is managed for pulpwood production by clear-cutting, followed by site preparation and direct seeding of sand pine. Recent research indicates only minor differences in vegetation recov-


Table 1. Preburn shrub cover and fire intensity for prescribed burn of mature Ocala sand pine scrub.

Plot	Location	Shrub cover (%)	Mean intensity
1	Edge	79	1.7
2	Interior	74	4.3
3	Edge	92	1.3
4	Interior	60	1.5
5	Edge	82	1.3
6	Interior	80	3.9


## RESULTS AND DISCUSSION

The burn was a successful operation. Although it produced a smoke column visible up to 25 kilometers away, the smoke rose quickly and dispersed rapidly creating no management problems (Custer and Thorsen 1996). Some minor spotting occurred during the backfire stage, but the spots were easily controlled. This study shows that high-intensity prescribed burns can be successfully implemented in the interior sand pine scrub of central Florida with careful planning.

Fire intensity varied with location and fuel loading



PLOT 4



PLOT 5




Figure 4. Relative intensity of fire across plots 4 and 5 (0 = no damage, 1 = tag tarnished, 2 = lower end of tag slightly melted, 3 = lower half of tag melted, 4 = 75% of the tag melted, 5 = tag completely melted, 6 = tag and nail melted, 7 = tag and nail burned completely off).

within the burned area (Table 1). Intensity was lower in plots 1 and 3 near the edge and increased toward the northwest as the fire gained strength (Figure 3). Most of plot 5 (Figure 4) was burned with a low-intensity backfire. Greatest fire intensities were in the interior of the burned area as shown by data from plots 2 and 6 (Figure 5). Plot 4 was also in the interior, but did not burn as intensely as expected (Figure 4). Most of plot 4 was on the middle portion of a slope where conditions were drier and the sparse woody understory covered only 60% of the plot (Table 1). This lower fuel loading resulted in the lower observed intensity. Carrington (1996) also reported high variability of fire temperature at the soil surface during the burn.

Before prescribed burning the stand had  $670 \pm 54$  Ocala sand pines/hectare,  $77 \pm 16$  oaks/hectare and  $137 \pm 30$  standing snags/hectare in the overstory. Sand pine diameter ranged from 6 to 30 centimeters with an average of  $17.5 \pm 0.3$  centimeters and a mean height of  $16.8 \pm 0.4$  meters. The oaks had an average diameter of  $6.4 \pm 0.6$  centimeters and a mean height of  $4.2 \pm 0.3$  meters. The fire felled 27% of the preexisting snags and 4% of the living oaks greater than 6

Figure 3. Relative intensity of fire across plots 1 and 3 (0 = no damage, 1 = tag tarnished, 2 = lower end of tag slightly melted, 3 = lower half of tag melted, 4 = 75% of the tag melted, 5 = tag completely melted, 6 = tag and nail melted, 7 = tag and nail burned completely off).

Death of the overstory, creation of new snags, elimination of the understory, and reduction in the litter layer significantly changed the structure of this patch of scrub. The plant community has been successfully reset to the beginning point where early successional plant and animal species requiring open scrub habitat can again thrive.

## ACKNOWLEDGMENTS

We thank the staff of the Seminole Ranger District, Ocala National Forest, U.S. Department of Agriculture, Forest Service, for conducting the prescribed burn. Dr. Tom Waldrop and Dr. Tom Lloyd provided valuable reviews for manuscript improvement. We are indebted to Patricia A. Outcalt and Liz O'Neill for help in study installation and data collection.

## LITERATURE CITED

- Brendemuehl, R.H. 1990. *Pinus clausa* (Chapm. ex Engelm.) Vasey ex Sarg. Sand Pine. Pages 294–301 in R.M. Burns and B.H. Honkala (technical coordinators). *Silvics of North America: 1. Conifers*. Agriculture Handbook 654. U.S. Department of Agriculture, Forest Service, Washington, DC.
- Brooks, H.K. 1972. The geology of the Ocala National Forest. Pages 81–92 in S.C. Snedaker and A.E. Lugo (eds.). *Ecology of the Ocala National Forest*. U.S. Department of Agriculture, Forest Service, Southeast Region, Atlanta, Georgia.
- Carrington, M.E. 1996. Post fire recruitment in Florida sand pine scrub in comparison with California Chaparral. Dissertation. University of Florida, Gainesville.
- Christman, S.P., and W.S. Judd. 1990. Notes on plants endemic to Florida scrub. *Biological Science* 53:52–73.
- Custer, G., and J. Thorsen. 1996. Stand-replacement burn in the Ocala National Forest—A success. *Fire Management Notes* 56:7–12.
- Doren, R.F., D.R. Richardson, and R.E. Roberts. 1987. Prescribed burning of the sand pine scrub community: Yamato scrub, a test case. *Florida Scientist* 50:184–192.
- Greenberg, C.H., D.G. Neary, and L.D. Harris. 1994. Effect of high-intensity wildfire and silvicultural treatments on reptile communities in sand-pine scrub. *Conservation Biology* 8: 1047–1057.
- Greenberg, C.H., L.D. Harris, and D.G. Neary. 1995a. A comparison of bird communities in burned and salvage-logged, clear-cut, and forested Florida sand pine scrub. *Wilson Bulletin* 107:40–54.
- Greenberg, C.H., D.G. Neary, L.D. Harris, and S.P. Linda. 1995b. Vegetation recovery following high-intensity wildfire and silvicultural treatments in sand pine scrub. *American Midland Naturalist* 133:149–163.
- Harper, R.M. 1927. *Natural resources of southern Florida*. 18th Annual Report, Florida Geologic Survey, Tallahassee.
- Hartnett, D.C., and D.R. Richardson. 1989. Population biology of *Bonamia grandiflora* (Convolvulaceae) effects of fire on plant and seed dynamics. *American Journal of Botany* 76: 361–369.
- Johansen, R.W., and D.D. Wade. 1987. Effects of crown scorch on survival and diameter growth of slash pines. *Southern Journal of Applied Forestry* 11:180–184.
- Johnson, S.F. 1982. Some demographic characteristics of the Florida rosemary *Ceratiola ericoides* Michx. *American Midland Naturalist* 108:170–174.
- Keetch, J.J., and G.M. Byram. 1968. A drought index for forest fire control. Research Paper SE-38, U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station, Asheville, NC.
- Kurz, H. 1942. Florida dunes and scrub, vegetation and geology. Bulletin No. 23, Florida Geologic Survey, Tallahassee.
- Laessle, A.M. 1958. The origin and successional relationship of sandhill vegetation and sand pine scrub. *Ecological Monographs* 28:361–387.
- Myers, R.L. 1990. Scrub and high pine. Pages 150–193 in R.L. Myers and J.J. Ewel (eds.). *Ecosystems of Florida*. University of Central Florida Press, Orlando.
- Small, J.K. 1921. Old trails and new discoveries. A record of exploration in Florida in the spring of 1919. *Journal of The New York Botanical Garden* 22:25–40; 49–64.
- Stocks, B.J. 1989. Fire behavior in mature jack pine. *Canadian Journal of Forest Research* 19:783–790.
- Veno, P.A. 1976. Successional relationships of five Florida plant communities. *Ecology* 57:498–508.
- Wade, D.D., and R.W. Johansen. 1987. Relating wildland fire to defoliation and mortality in pine. Pages 107–110 in D. R. Phillips (compiler). *Proceedings of the fourth biennial southern silvicultural research conference*. General Technical Report SE-42, U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station, Asheville, NC.